

Indications

For Monovision Treatment of Presbyopic Patients with Low to Moderate Nearsightedness (Myopia) with and without Astigmatism; myopia (nearsightedness) up to -11.00 D , with astigmatism up to -5.00 D; mixed astigmatism from 1 to 5 diopters; Hyperopia (Farsightedness) up to 4 Diopters (D) with Astigmatism, between 0 and 2 D

Important Information:

- **iDESIGN**® System driven LASIK Treatments is a permanent operation to the cornea and is irreversible.
- **iDESIGN**® System driven LASIK Treatments may not eliminate the need for reading glasses.
- Your vision must be stable for at least one year before **iDESIGN**® System driven LASIK Treatments. You will need written evidence of a change in manifest refraction of no more than +0.5 D (in both cylinder and sphere components) for at least one year prior to the date of pre-operative examination for Monovision LASIK; near sightedness, mixed astigmatism and farsightedness has not changed less than 1.0 dioter for respective indication.
- Pregnant and nursing women should wait until they are not nursing and not pregnant to have the surgery.
- You are not a good candidate if you have degenerative or autoimmune diseases, or have a condition that makes wound healing difficult.
- **iDESIGN**® System driven LASIK Treatments may result in some discomfort. The surgery is not risk-free. Please read patient information booklet, especially the sections on Benefits and Risks before you agree to the surgery.
- Alternatives to **iDESIGN**® System driven LASIK Treatments, but are not limited to, glasses, contact lenses, RK, and ALK.
- **iDESIGN**® System driven LASIK Treatments cannot meet the job-related vision requirements for some people such as pilots.
- You should be willing to undergo a one-week contact lens trial with your individualized monovision prescription and evaluate your vision over a range of tasks during this trial period, this will help you find out if you can tolerate Monovision LASIK.
- It may take several weeks for your eyes to adjust to monovision. You may want to avoid or decrease your participation in visually demanding situations such as driving until you have adjusted to your monovision.
- It is very important to stop wearing contact lenses before the pre-surgical eye exam.
- Before considering laser vision correction you should:
 - a. Have a complete eye exam.
 - b. Talk with one or more eye care professionals about the potential benefits of laser refractive surgery, and the complications, risks, and time required for healing.